


Q1 In which geographic region(s) are you licensed? Please check all that apply:


Answered: 198 Skipped: 3



Answer Choices	Percentage	Responses
● East Asia and Pacific	4.98%	10
● Europe and Central Asia	29.35%	59
● Latin America and Caribbean	8.96%	18
● Middle East and North Africa	1.99%	4
● North America	51.24%	103
● South Asia	2.49%	5
● Sub-Saharan Africa	1.00%	2
Total		201

Q2 I sedate and/or anesthetize birds presented to my practice for euthanasia.


Answered: 199 Skipped: 2



Answer Choices	Percentage	Responses
● Never	3.02%	6
● Rarely: Less than 10% of the time	9.05%	18
● Occasionally: About 30% of the time	4.02%	8
● Sometimes: About 50% of the time	4.52%	9
● Frequently: About 70% of the time	16.58%	33
● Always	62.81%	125
Total		199

Q3 I sedate and/or anesthetize psittacine birds presented for euthanasia.


Answered: 200 Skipped: 1



Answer Choices	Percentage	Responses
● Never	5.00%	10
● Rarely: Less than 10% of the time	5.00%	10
● Occasionally: About 30% of the time	2.00%	4
● Sometimes: About 50% of the time	4.50%	9
● Frequently: About 70% of the time	9.50%	19
● Always	74.00%	148
Total		200

Q4 I prescribe pre-visit anxiety relief for companion birds that will be presented for euthanasia.


Answered: 196 Skipped: 5



Answer Choices	Percentage	Responses
● Never	49.49%	97
● Rarely: Less than 10% of the time	33.16%	65
● Occasionally: About 30% of the time	8.67%	17
● Sometimes: About 50% of the time	5.10%	10
● Frequently: About 70% of the time	1.53%	3
● Always	2.04%	4
Total		196

Q5 For sedation/anesthesia of birds that will be euthanized, I commonly use the follow routes of administration (SELECT ALL THAT APPLY):


Answered: 191 Skipped: 10



Answer Choices	Percentage	Responses
● Oral/transmucosal	7.99%	31
● Inhalation	35.05%	136
● Intranasal	20.62%	80
● Parenteral	36.34%	141
Total		388

Q6 I routinely place an IV catheter in companion birds that will be euthanized in the presence of an owner (when patient size allows)


Answered: 195 Skipped: 6



Answer Choices	Percentage	Responses
● Never	36.92%	72
● Rarely: Less than 10% of the time	25.64%	50
● Occasionally: About 30% of the time	5.13%	10
● Sometimes: About 50% of the time	10.77%	21
● Frequently: About 70% of the time	9.74%	19
● Always	11.79%	23
Total		195

Q7 I regularly use the following inhaled agents to euthanize birds (SELECT ALL THAT APPLY):


Answered: 198 Skipped: 3



Answer Choices	Percentage	Responses
● Inhaled anesthetics	36.19%	76
● Argon noble gas (Ar)	0%	0
● Carbon dioxide (CO2)	2.38%	5
● Carbon monoxide (CO)	0.48%	1
● Nitrogen (N2)	0%	0
● I do not regularly use inhalant agents to euthanize birds.	57.62%	121
● Other (please specify) Show responses	3.33%	7
Total		210

Q8 I regularly use the following injectable agents to euthanize birds (SELECT ALL THAT APPLY):


Answered: 196 Skipped: 5



Answer Choices	Percentage	Responses
● Barbiturates	74.45%	169
● Potassium chloride	9.25%	21
● T-61	8.81%	20
● I do not regularly use injectable agents to euthanize birds.	3.08%	7
● Other (please specify) Show responses	4.41%	10
Total		227

Q9 When administering injectable euthanasia agents to birds, I regularly use the following route(s) of administration (SELECT ALL THAT APPLY):


Answered: 194 Skipped: 7



Answer Choices	Percentage	Responses
● Intracardiac	25.94%	117
● Intracoelomic	12.20%	55
● Intrahepatic	10.64%	48
● Intravenous, peripheral vein	40.13%	181
● Intravenous, occipital venous sinus	7.98%	36
● Oral transmucosal	3.10%	14
Total		451

Q10 My preferred route for administration of injectable euthanasia agents given to companion birds is... Rank your preferred route first, then list any other routes used in descending order. Leave blank if not used.


Answered: 191 Skipped: 10



	1	2	3	4	5	6	TOTAL	SCORE
Intracardiac	14.55% 24	44.85% 74	21.21% 35	13.33% 22	4.85% 8	1.21% 2	165	4.47
Intracoelemic	5.69% 7	19.51% 24	29.27% 36	26.83% 33	15.45% 19	3.25% 4	123	3.63
Intrahepatic	3.28% 4	17.21% 21	34.43% 42	24.59% 30	14.75% 18	5.74% 7	122	3.52
Intravenous, peripheral vein	77.66% 146	12.23% 23	2.66% 5	4.26% 8	0.53% 1	2.66% 5	188	5.54
Intravenous, occipital venous sinus	2.75% 3	22.02% 24	16.51% 18	19.27% 21	24.77% 27	14.68% 16	109	3.15
Oral transmucosal	2.20% 2	9.89% 9	8.79% 8	4.40% 4	17.58% 16	57.14% 52	91	2.03

Q11 My preferred route for administration of injectable euthanasia agents given to aviary birds is... (Rank your preferred route first, then list any other routes used in descending order. Leave blank if not used):


Answered: 174 Skipped: 27



	1	2	3	4	5	6	TOTAL	SCORE
Intracardiac	20.28% 29	41.26% 59	21.68% 31	13.99% 20	2.80% 4	0.00% 0	143	4.62
Intracoelomic	11.11% 12	21.30% 23	25.00% 27	25.93% 28	12.96% 14	3.70% 4	108	3.81
Intrahepatic	5.77% 6	25.96% 27	28.85% 30	25.00% 26	9.62% 10	4.81% 5	104	3.79
Intravenous, peripheral vein	71.07% 113	11.95% 19	6.92% 11	3.77% 6	2.52% 4	3.77% 6	159	5.34
Intravenous, occipital venous sinus	7.53% 7	22.58% 21	13.98% 13	10.75% 10	34.41% 32	10.75% 10	93	3.26
Oral transmucosal	3.90% 3	6.49% 5	10.39% 8	6.49% 5	11.69% 9	61.04% 47	77	2.01

Q12 My preferred route for administration of injectable euthanasia agents given to wild avian patients is... (Rank your preferred route first, then list any other routes used in descending order. Leave blank if not used):


Answered: 186 Skipped: 15



	1	2	3	4	5	6	TOTAL	SCORE
Intracardiac	18.71% 29	43.87% 68	23.23% 36	10.32% 16	3.87% 6	0.00% 0	155	4.63
Intracoelemic	7.63% 9	23.73% 28	27.12% 32	29.66% 35	8.47% 10	3.39% 4	118	3.82
Intrahepatic	2.61% 3	22.61% 26	31.30% 36	28.70% 33	12.17% 14	2.61% 3	115	3.67
Intravenous, peripheral vein	72.09% 124	11.63% 20	6.40% 11	2.91% 5	3.49% 6	3.49% 6	172	5.35
Intravenous, occipital venous sinus	12.77% 12	20.21% 19	13.83% 13	10.64% 10	34.04% 32	8.51% 8	94	3.41
Oral transmucosal	2.67% 2	4.00% 3	5.33% 4	5.33% 4	13.33% 10	69.33% 52	75	1.69

Q13 My preferred route for administration of injectable euthanasia agents given to pediatric avian patients is ... (Rank your preferred route first, then list any other routes used in descending order. Leave blank if not used):

Answered: 173 Skipped: 28



	1	2	3	4	5	6	TOTAL	SCORE
Intracardiac	37.33% 56	37.33% 56	16.00% 24	7.33% 11	2.00% 3	0.00% 0	150	5.01
Intracoelemic	17.95% 21	23.08% 27	25.64% 30	23.93% 28	5.98% 7	3.42% 4	117	4.13
Intrahepatic	13.76% 15	21.10% 23	37.61% 41	13.76% 15	11.93% 13	1.83% 2	109	4.06
Intravenous, peripheral vein	45.99% 63	14.60% 20	12.41% 17	14.60% 20	8.03% 11	4.38% 6	137	4.63
Intravenous, occipital venous sinus	10.71% 9	17.86% 15	9.52% 8	14.29% 12	34.52% 29	13.10% 11	84	3.17
Oral transmucosal	6.67% 5	8.00% 6	5.33% 4	9.33% 7	10.67% 8	60.00% 45	75	2.11